(Ohba, Kikuchi, Wakabayashi, Suzuki, Kurosaki, Rajbhandari & Wu 8570278, TI, holotype, KATH, isotype); Dubikharka (3720 m)-Baula Pokhari (3960 m), 9 July 1985, in forest (Ohba, Kikuchi, Wakabayashi, Suzuki, Kurosaki, Rajbhandari & Wu 8570329, TI, KATH, BM, NY).

We are indebted to Dr. S. B. Malla, Director General and Dr. S. B. Rajbhandari, Deputy Director General, H. M. G. Department of Medicinal Plants, Kathmandu, Nepal for providing us opportunities to study the Himalayan plants.

References

Clarke, C. B. 1894. Cyperaceae. *In*: J. D. Hooker, Flora of British India **6**: 585–748. Koyama, T. 1978. Cyperaceae. *In*: H. Hara et al., An enumeration of the flowering plants of Nepal 1: 96–120. Kükenthal, G. 1909. Cyperaceae-Caricoideae. *In*: A. Engler, Pflanzenreich IV-20, Ht. 38.

* * * *

北半球の温帯に分布するヒゲハリスゲ属は，中央アジアおよびヒマラヤで特に多様化している。ヒゲハリスゲ属植物は，ヒマラヤの高山帯の植物相の主要な構成種であるにもかかわらず，その全体像は十分に把握されていない。ここで記載した新種は，東ネパールのラメャップ地域の亜高山帯の森林内で採取されたもので，近似する *Kobresia Williamsii* から，植物体の高さ，前葉の大きさ，小穂の苞えいの先が丸いことにより区別される。

□Nishida, M. (ed.): **Contributions to the botany in the Andes II** 158 pp. 66 pls. 1987. アカデミア洋書，東京. ¥11,500. 千葉大学の西田誠教授を中心にすすめられている南米のアンデスを中心とする地域の植物調査の報告書である。この調査は「文部省海外学術調査」によるもので，チリの Quiriquina 島の上部白亜紀化石材の研究（西田誠・西田治文），アンデス山脈の東西両斜面に発達する雲霧林の生態（1. 微気象，及び 2. チリ中南部の *Nothofagus* 林の群落構造；共に大賀宣彦），南アメリカ産シダ植物の染色体研究（栗田子郎），及びチリ産の重要樹種の材組織の研究（M. Rancusi, H.他）の5編の論文からなる。なお，本調査報告書のIは1985年に千葉大学理学部から発行されている。 （井上 浩）